

Bayer® Back & Body

Drug Facts

Active ingredients (in each caplet)	Purpose
Aspirin 500 mg (NSAID)*.....	Pain reliever
Caffeine 32.5 mg.....	Pain reliever aid

*nonsteroidal anti-inflammatory drug

Uses for the temporary relief of

- backache pain • muscle aches and pains
- minor aches and pains of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives • facial swelling • asthma (wheezing) • shock
- Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you
- are age 60 or older
 - have had stomach ulcers or bleeding problems
 - take a blood thinning (anticoagulant) or steroid drug
 - take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
 - have 3 or more alcoholic drinks every day while using this product
 - take more or for a longer time than directed

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout • diabetes • arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint • vomit blood • have bloody or black stools
 - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

If pregnant or breast-feeding, ask a health professional before use.

It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.



Bayer® Back & Body

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 2 caplets every 6 hours not to exceed 8 caplets in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients carnauba wax, corn starch, D&C red #30 aluminum lake, D&C yellow #10 aluminum lake, hypromellose, powdered cellulose, propylene glycol, shellac, titanium dioxide, triacetin

Questions or comments? 1-800-331-4536 (Mon-Fri 9AM - 5PM EST)