

## Aspirin Regimen

### Bayer® Chewable Low Dose Aspirin 81 mg Cherry Flavored

#### Drug Facts

Active ingredient (in each tablet)	Purpose
Aspirin 81 mg (NSAID)*.....	Pain reliever
*nonsteroidal anti-inflammatory drug	

#### Uses

- for the temporary relief of minor aches and pains or as recommended by your doctor
- ask your doctor about other uses for Bayer 81 mg Cherry Chewable Aspirin

#### Warnings

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives • facial swelling • asthma (wheezing) • shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

#### Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- this product for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients

#### Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

**Ask a doctor or pharmacist before use if you are taking a prescription drug for**

- gout • diabetes • arthritis

#### Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
  - feel faint • vomit blood • have bloody or black stools
  - have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- redness or swelling is present
- new symptoms occur
- ringing in the ears or a loss of hearing occurs

**If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.



## Aspirin Regimen

### Bayer® Chewable Low Dose Aspirin 81 mg Cherry Flavored

**Directions**

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

**Other information**

- save carton for full directions and warnings
- store at room temperature

**Inactive ingredients** colloidal silicon dioxide, corn starch, D&C red #27 aluminum lake, dextrose, FD&C red #40 aluminum lake, flavor, microcrystalline cellulose, saccharin sodium

**Questions or comments? 1-800-331-4536 (Mon-Fri 9AM – 5PM EST)**