

Aspirin Regimen

Bayer® Chewable Low Dose Aspirin 81 mg Orange Flavored

Drug Facts	
Active ingredient (in each chewable tablet)	Purpose
Aspirin 81 mg (NSAID)*.....	Pain reliever
*nonsteroidal anti-inflammatory drug	
Uses	
<ul style="list-style-type: none"> • for the temporary relief of minor aches and pains or as recommended by your doctor • ask your doctor about other uses for Bayer 81 mg Orange Chewable Aspirin 	
Warnings	
<p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction which may include:</p> <ul style="list-style-type: none"> • hives • facial swelling • asthma (wheezing) • shock <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you</p> <ul style="list-style-type: none"> • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed 	
Do not use	
<ul style="list-style-type: none"> • if you are allergic to aspirin or any other pain reliever/fever reducer • this product for at least 7 days after tonsillectomy or oral surgery unless directed by a doctor • if you have ever had an allergic reaction to this product or any of its ingredients 	
Ask a doctor before use if	
<ul style="list-style-type: none"> • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma 	
Ask a doctor or pharmacist before use if you are taking a prescription drug for	
<ul style="list-style-type: none"> • gout • diabetes • arthritis 	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> • an allergic reaction occurs. Seek medical help right away. • you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • ringing in the ears or a loss of hearing occurs 	
<p>If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.</p> <p>Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.</p>	



Aspirin Regimen
Bayer® Chewable Low Dose Aspirin 81 mg
Orange Flavored

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 4 to 8 tablets every 4 hours not to exceed 48 tablets in 24 hours unless directed by a doctor
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients colloidal silicon dioxide, corn starch, dextrose, FD&C yellow #6 aluminum lake, flavor, microcrystalline cellulose, saccharin sodium

Questions or comments? 1-800-331-4536 (Mon-Fri 9AM – 5PM EST)