



Claritin RediTabs 24 Hour

Drug Facts

Active ingredient (in each tablet)	Purpose
Loratadine 10 mg.....	Antihistamine

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- place 1 tablet on tongue; tablet disintegrates, with or without water

adults and children 6 years and over	1 tablet daily; not more than 1 tablet in 24 hours
--------------------------------------	--

children under 6 years of age	ask a doctor
-------------------------------	--------------

consumers with liver or kidney disease	ask a doctor
--	--------------

Other information

- safety sealed: do not use if the individual blister unit imprinted with Claritin® RediTabs® is open or torn
- store between 20° to 25° C (68° to 77° F)
- use tablet immediately after opening individual blister
- complies with USP Procedure 2 for Assay and Organic Impurities and Test 2 for Disintegration

Inactive ingredients

anhydrous citric acid, gelatin, mannitol, mint flavor

Questions or comments?

1-800-CLARITIN (1-800-252-7484) or www.claritin.com