

Drug Facts

Active ingredient (in each tablet)

Loratadine 10 mg..... Antihistamine

Purpose

Uses temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:

- runny nose
- itchy, watery eyes
- sneezing
- itching of the nose or throat

Warnings

Do not use if you have ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if you have liver or kidney disease. Your doctor should determine if you need a different dose.

When using this product do not take more than directed. Taking more than directed may cause drowsiness.

Stop use and ask a doctor if an allergic reaction to this product occurs. Seek medical help right away.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- chew or crush tablets completely before swallowing.

adults and children 6 years and over	chew 1 tablet daily; not more than 1 tablet in 24 hours
children under 6 years of age	ask a doctor
consumers with liver or kidney disease	ask a doctor

Other information

- phenylketonurics: contains phenylalanine 3 mg per tablet
- safety sealed: do not use if the individual blister unit imprinted with Claritin® is open or torn
- store between 20° to 25°C (68° to 77°F)

Inactive ingredients

aspartame, citric acid anhydrous, colloidal silicon dioxide, FD&C blue No. 1 aluminum lake, flavors, magnesium stearate, mannitol, menthol, microcrystalline cellulose, sodium starch glycolate, stearic acid

Questions or comments?

1-800-CLARITIN (1-800-252-7484) or www.claritin.com

RESTRICTED