



Genuine Bayer® Aspirin Tablets

Drug Facts	
Active ingredient (in each tablet)	Purposes
Aspirin 325 mg (NSAID)*.....	Pain reliever/fever reducer
*nonsteroidal anti-inflammatory drug	
Uses	
<ul style="list-style-type: none"> • temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> • headache • muscle pain • toothache • menstrual pain • colds • minor pain of arthritis • temporarily reduces fever 	
Warnings	
<p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.</p> <p>Allergy alert: Aspirin may cause a severe allergic reaction which may include:</p> <ul style="list-style-type: none"> • hives • facial swelling • asthma (wheezing) • shock <p>Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you</p> <ul style="list-style-type: none"> • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while using this product • take more or for a longer time than directed 	
Do not use	
<ul style="list-style-type: none"> • if you are allergic to aspirin or any other pain reliever/fever reducer • if you have ever had an allergic reaction to this product or any of its ingredients 	
Ask a doctor before use if	
<ul style="list-style-type: none"> • stomach bleeding warning applies to you • you have a history of stomach problems, such as heartburn • you have high blood pressure, heart disease, liver cirrhosis, or kidney disease • you are taking a diuretic • you have asthma 	
Ask a doctor or pharmacist before use if you are taking a prescription drug for	
<ul style="list-style-type: none"> • gout • diabetes • arthritis 	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> • an allergic reaction occurs. Seek medical help right away. • you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • ringing in the ears or a loss of hearing occurs 	



Genuine Bayer® Aspirin Tablets

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.** **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose
- adults and children 12 years and over: take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours
- children under 12 years: consult a doctor

Other information

- save carton for full directions and warnings
- store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients corn starch, hypromellose, powdered cellulose, triacetin

Questions or comments? 1-800-331-4536 (Mon – Fri 9AM – 5PM EST)