



Coricidin HBP

Chest Congestion & Cough

Drug Facts

Active ingredients (in each softgel)	Purpose
Dextromethorphan hydrobromide 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant

Uses

- temporarily relieves cough due to minor throat and bronchial irritation as may occur with a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to make coughs more productive

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- cough that occurs with excessive phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if cough lasts more than 7 days, reoccurs, or occurs with fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not exceed recommended dose
- adults and children 12 years and over: take 1 or 2 softgels with water every 4 hours. Do not exceed 12 softgels in 24 hours or as directed by a doctor.
- children under 12 years of age: do not use

Other information

- store between 20° to 25°C (68° to 77°F)
- protect from freezing
- protect from excessive moisture

Inactive ingredients FD&C blue no.1, FD&C red no. 40, gelatin, glycerin, pharmaceutical ink, polyethylene glycol 400, povidone, propylene glycol, purified water, sorbitol

Questions or comments? 1-800-317-2165 (Mon-Fri 9AM - 5PM EST)