



Alka-Seltzer Plus[®] Maximum Strength Day Cold & Flu Liquid Gels

| Drug Facts | |
|---|-----------------------------|
| Active ingredients (in each capsule) | Purposes |
| Acetaminophen 325 mg..... | Pain reliever/fever reducer |
| Dextromethorphan hydrobromide 10 mg..... | Cough suppressant |
| Phenylephrine hydrochloride 5 mg..... | Nasal decongestant |
| Uses | |
| <ul style="list-style-type: none">temporarily relieves these symptoms due to a cold or flu:<ul style="list-style-type: none">minor aches and painsheadachecoughsore throatnasal and sinus congestiontemporarily reduces fever | |
| Warnings | |
| Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take <ul style="list-style-type: none">more than 4,000 mg of acetaminophen in 24 hourswith other drugs containing acetaminophen3 or more alcoholic drinks every day while using this product | |
| Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: <ul style="list-style-type: none">skin reddeningblistersrashhivesfacial swellingasthma (wheezing)shock If a skin or general allergic reaction occurs, stop use and seek medical help right away. | |
| Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. | |
| Do not use | |
| <ul style="list-style-type: none">with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.if you have ever had an allergic reaction to this product or any of its ingredientsin children under 12 years of age | |
| Ask a doctor before use if you have | |
| <ul style="list-style-type: none">liver diseaseheart diseasehigh blood pressurethyroid diseasediabetescough with excessive phlegm (mucus)difficulty in urination due to enlargement of the prostate glandpersistent or chronic cough such as occurs with smoking, asthma, or emphysema | |
| Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin | |
| When using this product do not exceed recommended dosage | |
| Stop use and ask a doctor if | |
| <ul style="list-style-type: none">pain, cough, or nasal congestion gets worse or lasts more than 7 daysfever gets worse or lasts more than 3 daysredness or swelling is presentnew symptoms occurcough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.nervousness, dizziness, or sleeplessness occurs | |



Alka-Seltzer Plus® Maximum Strength Day Cold & Flu Liquid Gels

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years: do not use

Other information

- store at room temperature. Avoid excessive heat above 40°C (104°F).

Inactive ingredients FD&C yellow #6, FD&C red #40, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

Questions or comments? 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)